

Dark Matter, Dark Energy and the Chaplygin Gas

Neven Bilić[†], Gary B. Tupper, and Raoul D. Viollier[‡]
 Institute of Theoretical Physics and Astrophysics,
 Department of Physics, University of Cape Town,
 Private Bag, Rondebosch 7701, South Africa

February 1, 2008

Abstract

We formulate a Zel'dovich-like approximation for the Chaplygin gas equation of state $P = -A/\rho$, and sketch how this model unifies dark matter with dark energy in a geometric setting reminiscent of M -theory.

1 Introduction

In the last few years improved observations [1] have forced a shift in our cosmological paradigm: the $\Omega_M = 1$ dust model has been swept aside and, in its place, we are faced with the problem of understanding a universe with an equation of state $\bar{W} = \bar{P}/\bar{\rho} < -1/3$. That is to say, on average, pressure is comparable with density and, moreover, negative.

Of course, parametrically, this is readily accommodated by a cosmological constant Λ [2] with $\Omega_\Lambda \simeq 0.7$ and $\Omega_{DM} = 1 - \Omega_\Lambda \simeq 0.3$ (throughout the paper we neglect the small baryonic contribution). The well-known difficulty with Λ is that *a priori* it seems an incredible accident that $\Omega_\Lambda \simeq \Omega_{DM}$ since

[†]Permanent address: Rudjer Bošković Institute, P.O. Box 180, 10002 Zagreb, Croatia;
 Email: bilic@thphys.irb.hr

[‡]Email: viollier@physci.uct.ac.za

$\rho_\Lambda/\rho_{\text{DM}} \sim a^3$, a being the scale factor. Hence, much attention has been devoted to quintessence [3], involving a real scalar field which tracks [4] the background component until recently becoming dominant. However, simple tracking quintessence does not work [5] and spintessence [6], where the scalar field is complex, suffers instabilities against the decay of dark energy into dark matter [7].

It is natural to conjecture that some of the aforementioned problems derive from treating dark matter and dark energy as separate issues. As an example, Barr and Seckel [8] have pointed out that in axion dark matter models quantum gravity effects break the Pecci-Quinn symmetry leading to a universe trapped in a false vacuum with an effective Λ of the correct magnitude. In another approach, Wetterich [9] has suggested that traditional WIMP dark matter should be replaced by quintessence lumps, thus unifying dark matter and dark energy. However, the radiation-matter transition and structure formation remain open questions in this scenario.

Herein we present a dark matter-energy unification model suggested by the observation of Kamenshchik et al. [10] that a perfect fluid obeying the Chaplygin gas equation of state

$$P = -\frac{A}{\rho} \tag{1}$$

should lead to a homogenous cosmology with

$$\bar{\rho}(a) = \sqrt{A + \frac{B}{a^6}}, \tag{2}$$

with B being an integration constant, thus interpolating between dark matter, $\bar{\rho}(a \rightarrow 0) \simeq \sqrt{B}/a^3$ and dark energy $\bar{\rho}(a \rightarrow \infty) \simeq \sqrt{A}$. Before doing so, we must first show why Eq. (1), aside from its interesting mathematical features [11], might describe reality.

2 Brane New World

One of the most profound recent developments in fundamental physics has been the recognition that all of the extra dimensions required by string/ M -theory do not have to be of the Planck length size: one (or more) could be as large as 0.1 mm provided that all standard-model fields except gravity are

confined to a 3-dimensional hypersurface or ‘brane’ in the higher dimensional bulk (for a review see [12]). In this context, Kamenshchik et al. [13] obtained Eq. (1) from the stabilization of branes in black hole bulks.

A simple way to see the connection between the Chaplygin gas and the brane world is to follow Sundrum’s [14] effective field theory for the 3- brane. The gauge fixed embedding of a 3+1 brane in a 4+1 bulk is described by $Y^M = (x^\mu, Y^4)$. With some nominal assumptions on the bulk metric G_{MN} , the induced metric on the 3-brane is

$$\tilde{g}_{\mu\nu} = g_{\mu\nu} - \theta_{,\mu}\theta_{,\nu}, \quad (3)$$

with $0 \leq \theta \leq \ell_5$, and the action for the brane reads

$$\begin{aligned} S_{\text{brane}} &= \int d^4x \sqrt{-\tilde{g}} [-f^4 + \dots] \\ &= \int d^4x \sqrt{-g} \sqrt{1 - g^{\mu\nu} \theta_{,\mu} \theta_{,\nu}} [-f^4 - \dots], \end{aligned} \quad (4)$$

where f^4 is the brane tension and the ellipsis includes standard-model fields as well as higher-order terms in power counting. One estimates $f \sim \ell_5^{-1} \sim \text{meV}$.

Retaining only the leading term in Eq. (4) and renaming $f^4 = \sqrt{A}$, one sees that its content is equivalent to

$$\mathcal{L} = \frac{\phi^2}{2} g^{\mu\nu} \theta_{,\mu} \theta_{,\nu} - V \left(\frac{\phi^2}{2} \right), \quad (5)$$

$$V = \frac{1}{2} \left(\phi^2 + \frac{A}{\phi^2} \right) \quad (6)$$

since ϕ can be eliminated through its field equation

$$g^{\mu\nu} \theta_{,\mu} \theta_{,\nu} = V'. \quad (7)$$

We observe that $\mathcal{L}$ corresponds to the Lagrangian for a complex field $\Phi = \phi e^{-im\theta}/\sqrt{2m}$ in the ‘Thomas-Fermi’ approximation. The Thomas-Fermi approximation amounts to neglecting $\phi_{,\mu}/m\phi$ compared to $\sqrt{V'}/\phi^2$, i.e., the scale of variation of ϕ is large compared to the Compton wavelength. It is also worth noting that dividing $\mathcal{L}$ by $\sqrt{A}$, the first term is the periodic Gaussian model with coupling $R = A^{1/4}/\phi$. The potential $F = (R^2 + R^{-2})/2$, which is self-dual, can be interpreted as the mean field free-energy for ‘brane

cells' filling a system of size R , in analogy to Rama's [15] 'string bit' analysis of black holes.

To complete the connection to the Chaplygin gas, we point out a field-fluid duality: for $V' > 0$, Eq. (7) defines a fluid 4-velocity $U_\mu U^\mu = 1$,

$$U^\mu = g^{\mu\nu} \theta_{,\nu} / \sqrt{V'} \quad (8)$$

and then the energy-momentum tensor derived from $\mathcal{L}$ takes the perfect fluid form with

$$\rho = \frac{\phi^2}{2} V' + V, \quad P = \frac{\phi^2}{2} V' - V. \quad (9)$$

In particular, for V of Eq. (6), the equation of state (1) follows, and the energy density ρ is given by

$$\rho = \phi^2 = \frac{\sqrt{A}}{\sqrt{1 - g^{\mu\nu} \theta_{,\mu} \theta_{,\nu}}}, \quad (10)$$

which is to say that matter corresponds to a wrinkled brane.

Finally, it must be said that the procedure can be reversed [16]. The equations

$$d \ln(\phi^2) = \frac{d\rho - dP}{\rho + P}, \quad V = \frac{1}{2}(\rho - P), \quad (11)$$

obtained from Eq. (9), allow one to construct ϕ^2 and V given the equation of state. As an example, starting from $0 \geq W = P/\rho \geq -1$ and $0 \leq c_s^2 = dP/d\rho \leq 1$, with the relativistic limits coinciding, Eqs. (1), (6), and (4) follow.

3 The Inhomogeneous Chaplygin Gas

As yet, we have not dealt with the θ field equation; in the fluid language, it reads

$$\left(\sqrt{-g\phi^2(\rho)(\rho + P)} U^\mu \right)_{,\mu} = 0. \quad (12)$$

In comoving coordinates, $U^\mu = (1/\sqrt{g_{00}}, \vec{0})$, the solution for the Chaplygin gas is [16]

$$\rho = \sqrt{A + \frac{B}{\gamma}}. \quad (13)$$

Here $\gamma = -g/g_{00}$ is the determinant of the induced metric $\gamma_{ij} = g_{i0} g_{j0}/g_{00} - g_{ij}$ which measures physical distances, and $B = B(\vec{x})$ can be taken as constant on the scales of interest.

The generalization (13) of Eq. (2) allows us to implement the geometric version [17] of the Zel'dovich approximation [18]: the transformation from Lagrange to Euler (comoving) coordinates induces γ_{ij} as

$$\gamma_{ij} = \delta_{kl} D_i^k D_j^l, \quad D_i^j = a(\delta_i^j - b\varphi_{,i}^j), \quad (14)$$

with D_i^j the deformation tensor, φ the velocity potential. Inserting this ansatz in the 0-0 Einstein equation to first order in φ yields the evolution equation for $b(a)$

$$\frac{2}{3}a^2b'' + a(1 - \bar{w})b' = (1 + \bar{w})(1 - 3\bar{w})b, \quad (15)$$

$$\bar{w}(a) = -\frac{\Omega_\Lambda a^6}{1 - \Omega_\Lambda + \Omega_\Lambda a^6}, \quad (16)$$

where we match the parameters A, B to the Λ model.

In Fig. 1 we show the evolution of $b(a)$ for the Chaplygin gas and for Λ cold dark matter, the latter following by omitting the $(1 - 3\bar{w})$ factor in Eq. (15) and changing a^6 to a^3 in Eq. (16). In either case, the growth $b \propto a$ ceases near $a = 1$ and although b remains constant, the perturbative density contrast $\delta_{\text{pert}} = b(1 + \bar{w})\varphi_{,i}^i$ thereafter vanishes as $\delta_{\text{pert}}(a \gg 1) \sim a^{-6}$.

Of course the value of the Zel'dovich approximation is that it offers a means of extrapolation into the nonperturbative regime via Eqs. (13) and

$$\sqrt{\gamma} = a^3(1 - \lambda_1 b)(1 - \lambda_2 b)(1 - \lambda_3 b), \quad (17)$$

where the λ_i are the eigenvalues of $\varphi_{,i}^j$. When one (or more) of the λ 's is positive, a caustic forms on which $\gamma \rightarrow 0$ and $w \rightarrow 0$, i.e., at the locations where structure forms the Chaplygin gas behaves as dark matter. Conversely, when all of the λ 's are negative, a void forms, ρ is driven to its limiting value $\sqrt{A}$, and the Chaplygin gas behaves as dark energy driving accelerated expansion.

4 Discussion and Conclusions

A shortcoming of the Zel'dovich approximation is that at the caustic matter flows through unimpeded so that structures quickly dissolve [19]. This may




Figure 1: Evolution of $b(a)/b(a_{\text{eq}})$ from $a_{\text{eq}} = 1.0 \times 10^{-4}$ for $\Omega_{\Lambda} = 0.7$ and $b'(a_{\text{eq}}) = 0$, for the Chaplygin gas (solid line) and Λ CDM (dashed line).

be circumvented via the truncated Zel'dovich approximation [19]. A preferable alternative would be an extension of the adhesion approximation [20] which also allows the extraction of mass functions.

Approximation technicalities aside, the case is made that the Chaplygin gas offers a realistic unified model of dark matter and dark energy. That this is achieved in a geometric (brane world) setting rooted in string/ M theory makes this model all the more remarkable.

References

- [1] A. Melchiorri, these proceedings.
- [2] P.J.E. Peebles, ApJ **284**, 439 (1984); J.P. Ostriker and P.J. Steinhardt, Nature **377**, 600 (1995).
- [3] C. Wetterich, Nucl. Phys. B **302**, 668 (1988); P.J.E. Peebles and B. Ratra, ApJ **325**, L17 (1988); R.R. Caldwell, R. Dave and P.J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998).
- [4] I. Zlatev, L. Wang and P.J. Steinhardt, Phys. Rev. Lett. **82**, 896 (1999).
- [5] S. Bludman, these proceedings.
- [6] L.A. Boyle, R.R. Caldwell and M. Kamionkowski, astro-ph/0105318.
- [7] S. Kasuya, astro-ph/0105408.
- [8] S.M. Barr and D. Seckel, hep-ph/0106239.
- [9] C. Wetterich, hep-ph/0108266, astro-ph/0108411.
- [10] A. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B **511**, 265 (2001).
- [11] R. Jackiw, physics/0010042, and references therein.
- [12] V.A. Rubakov, hep-ph/0104152.
- [13] A. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B **487**, 7 (2000).
- [14] R. Sundrum, Phys. Rev. D **59**, 085009 (1999).
- [15] S.K. Rama, Phys. Lett. B **424**, 39 (1998).
- [16] N. Bilić, G.B. Tupper, and R.D. Viollier, astro-ph/0111325, Phys. Lett. B, in press.
- [17] S. Matarrese and D. Terranova, MNRAS **283**, 400 (1996).
- [18] Ya. B. Zel'dovich, A&A **5**, 84 (1970).
- [19] J.L. Pauls and A.L. Melott, MNRAS **274**, 99 (1995).

- [20] S.N. Gurbatov, A.I. Saichev and S.F. Shandarin, MNRAS **236**, 385 (1989).